**Accepted Manuscript (Uncorrected Proof)** 

Title: Nurses' Perspectives on Gender Equality Policies During the COVID-19 Response Phase

Authors: Safura Yaghmaei<sup>1</sup>, Abbasali Ebrahimian<sup>2,\*</sup>, Sima Pourteimour<sup>3</sup>

1. Department of Nursing, School of Nursing and Midwifery, Hamadan University of Medical

Sciences, Hamadan, Iran.

2. Disasters and Emergencies Group, Faculty of Paramedical Sciences, Qom University of Medical

Sciences, Qom, Iran.

3. Patient Safety Research Center, Clinical Research Institute, Urmia University of Medical Sciences,

Urmia, Iran.

To appear in: Health in Emergencies & Disasters Quarterly

**Received date: 2025/03/1** 

**Revised date: 2025/06/16** 

**Accepted date: 2025/07/29** 

First Online Published: 2025/07/29

This is a "Just Accepted" manuscript, which has been examined by the peer-review process and has been accepted for publication. A "Just Accepted" manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. *Health in Emergencies & Disasters Quarterly* provides "Just Accepted" as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the "Just Accepted" on Website and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

### Please cite this article as:

Yaghmaei S, Ebrahimian A, Pourteimour S. Nurses' Perspectives on Gender Equality Policies

During the COVID-19 Response Phase. Health in Emergencies & Disasters Quarterly.

Forthcoming 2026. DOI: http://dx.doi.org/10.32598/hdq.2026.141.2

DOI: http://dx.doi.org/10.32598/hdq.2026.141.2

### **Abstract:**

**Background:** Gender equality is a foundational aspect of the nursing profession. However, there is limited documentation of how gender considerations were incorporated into policymaking during the COVID-19 pandemic. This study explored nurses' perspectives on gender equality policies during the pandemic response phase.

**Materials and Methods:** A qualitative content analysis was conducted using semi-structured interviews with 14 nurses working in COVID-19 wards. Participants were selected through purposive sampling, continuing until data saturation. Data were analyzed following Graneheim and Lundman's method.

**Results:** The central theme identified was "gender inequality in management policies and resource allocation," derived from four categories: inefficient management, inequality in interactions, inequality in resource management, and inadequate attention to staff needs. Nurses reported experiencing gender bias in decision-making and observed unequal access to resources and support during the pandemic.

**Discussion:** Participants highlighted systemic gender disparities in management approaches during the crisis, including ineffective leadership and inequitable resource distribution. These findings are consistent with global evidence on gender-based inequalities during emergencies. To address these issues, the study recommends establishing gender-balanced leadership teams, implementing routine audits of resource distribution, and providing training in gender-sensitive management for healthcare administrators.

**Conclusion:** Nurses perceived significant gender inequality in policy-making during the COVID-19 response phase. Health system leaders and policymakers should prioritize gender equity in all stages of disaster preparedness and response. Further research is necessary to explore gender perspectives in similar high-stress healthcare environments.

Keywords: Nursing, Management, Policy making, Pandemic Response Plan, COVID-

### Introduction

A pandemic is defined as an epidemic that spreads across multiple countries or continents, affecting a large number of people. It represents a significant threat to global health, often leading to widespread morbidity and mortality, as well as social and economic disruptions. Historical examples include the Black Death and the Spanish flu, with contemporary instances such as COVID-19 (1, 2).

Nurses play a pivotal role in shaping pandemic response strategies through their frontline experiences, leadership, and adaptability. Their contributions are essential in developing frameworks that enhance healthcare systems' resilience during crises (3, 4). Nurses have developed structured response models, such as the 5A model (Alarm, Assessment, Adaptation, Amplification, Affect), which outlines their sequential responses to crises like COVID-19. This model serves as a theoretical foundation for future pandemic preparedness (3). Nurse leaders have demonstrated effective leadership by anticipating needs, fostering collaboration, and innovating care practices. Their experiences during the pandemic highlight the importance of involving nursing professionals in policy development and emergency preparedness planning (4).

The integration of gender mainstreaming into pandemic response strategies significantly enhances their effectiveness by addressing the unique vulnerabilities and needs of different genders, particularly women and girls. Gender mainstreaming ensures that pandemic responses are equitable and inclusive, which is crucial for mitigating the disproportionate impacts of health crises on women. This approach not only improves health outcomes but also supports broader social and economic stability during pandemics (5, 6). Gender-sensitive policies, such as those implemented in Kerala, India, have proven effective in addressing the specific needs of women during the COVID-19 pandemic. These policies included economic relief packages, support for mental health, and measures to combat gender-based violence, demonstrating the importance of gender considerations in policy-making (7). In contrast, the lack of gender-sensitive policies in Uganda led to increased gender inequalities, highlighting the need for proper gender analysis in designing response measures (5).

The presence of women in decision-making roles has been linked to more gender-sensitive policy responses. Countries with higher female representation in government taskforces, such as those led by women, tend to implement more inclusive and effective pandemic responses (8). However, women are often underrepresented in decision-making positions, which limits the integration of gender perspectives in pandemic planning and response (8, 9). An

intersectional approach to pandemic response, which considers the overlapping impacts of gender with other social determinants, is crucial for addressing the diverse needs of populations. This approach can help mitigate the compounded vulnerabilities faced by women, particularly those from marginalized communities (6, 10).

Despite the benefits of gender mainstreaming, many pandemic responses have failed to adequately incorporate gender perspectives, resulting in missed opportunities to address gender disparities effectively (11). The COVID-19 pandemic has highlighted the need for a genderresponsive, intersectional approach to future health crises, emphasizing the importance of integrating gender considerations into all levels of policy-making and implementation (10, 12). Given the intricacy and multidimensionality of gender justice, a comprehensive examination of this concept necessitates a thorough investigation within the relevant field. In this regard, qualitative studies of an in-depth nature offer a profound understanding of the subject matter under scrutiny. As such, the present research employs a qualitative approach to explore the nature of gender justice as experienced by nurses amid the COVID-19 pandemic. Previous research has merely dealt with gender justice in nursing and policymaking during various crises separately, and considering the continuation of the COVID-19 pandemic and the possibility of similar ones in the future, it is necessary to make policies based on gender equality from the perspectives of nurses working in COVID-19 units. Given the complexity and multidimensionality of concepts such as gender justice, a thorough investigation of said concepts within the desired field is imperative. In this regard, qualitative studies with an inherent depth are particularly valuable as they offer an intimate and insightful portrayal of the field under scrutiny. This study explored nurses' perspectives on gender equality policies during the response phase of the Covid-19 pandemic.

The unique contribution of this study lies in its focus on frontline nurses' perspectives regarding gender equality specifically within the context of management policies and resource allocation during the COVID-19 response phase. While previous research has broadly examined gender disparities in healthcare or workplace environments, this study highlights how these inequalities manifested in real-time crisis management—a context that exposed systemic gaps more starkly. This study breaks new ground by systematically examining structural gender inequities in crisis management during COVID-19, revealing how institutional policies perpetuated disparities despite formal equality measures. "Through in-depth qualitative analysis, this study uncovers the lived experiences of structural gender inequities in pandemic

crisis management, exposing how organizational cultures and hidden power dynamics systematically disadvantaged female healthcare nurses.

### **Methods**

### Study Design and Settings:

This study was conducted using a qualitative content analysis approach in 2020-2021. In this study, the Policy based on Gender Equality from the Perspective of Iranian Nurses in the COVID\_19 units was analyzed using content analysis through Graneheim and Lundman model (13). During the fieldwork phase, in-depth, face-to-face, and semi-structured interviews were conducted with 14 nurses who worked in COVID\_19 units, and each interview lasted between 30-min and 60-min. Participants were selected using purposive sampling. Data collection continued until reaching data saturation. All interviews were recorded. A total of 14 participants were selected through purposive sampling to ensure representation of diverse and relevant perspectives. This number was deemed sufficient for generating rich qualitative data within the scope of the study. Data saturation, the point at which no new themes or insights emerged, was used to determine the adequacy of the sample size. Saturation was reached by the 12th participant, and the final two interviews confirmed the consistency and completeness of the identified themes.

# Participants:

In selecting participants, maximum diversity was considered in terms of age, gender, and marital status (Table 1).

 Table 1. Profile of participants

participants	Age	Gender	Marital Status	experience
1	27	female	married	2 y
2	27.	male	single	3 y
3	35	male	married	10 y
4	30	female	single	6 y
5	32	male	single	8 y
6	30	male	married	5 y
7	28	male	married	4 y
8	34	female	single	10 y
9	37	female	married	14 y
10	35	male	married	9 y
11	30	female	single	6 y
12	37	female	married	12 y
13	33	female	single	10 y
14	29	female	single	6 y
Mean (SD)	31.71±3.49	Male 6 (42.85)	Married 7 (50)	7.5±3.5
or n(%)		Female 8 (57.14)	Single 7 (50)	

Inclusion criteria were at least two years of clinical work experience, Employment in one of the COVID\_19 units, Educational Hospitals and Medical Center in Iran, and Participant's desire to participate in the study. During the nurses' shift, the researcher referred to the nurses eligible to enter the study and placed those who wished to cooperate after the shift with them. The location and time of the interviews were selected according to the participants' preferences, but most interviews were conducted in restrooms after work shifts. Questions guiding interview at this phase included the following, but in the continuation of the interviews, with the progress of the study, data collection, simultaneous analysis, and formation of classes, the path of the subsequent interviews was determined:

- What were your biggest challenges at work during COVID-19?
- How do you see the effect of nurses 'gender differences in the COVID-19 pandemic on managers' decisions?
- Give an example of a policy that you found unfair or gender-based?
- What comes to mind when you come across the word gender justice in nursing policy?
- Was there a difference in access to protective equipment (e.g. masks, PPE) between male and female nurses?
- Did support policies (e.g., leave, psychological counseling) address the specific needs of female or male nurses?
- How do you see the performance of nursing managers in the COVID-19 crisis?

In addition, exploratory questions were also used to understand the interviewees' experiences better.

As the interviews progressed, questions were grouped by topic to guide the conversation and help explore various dimensions of the topic:

- 1. Background and Experience:
- A. Can you tell me about your role and responsibilities during the COVID-19 response phase?
  - B. How long have you been working in the nursing profession?
  - 2. Perceptions of Gender Equality in the Workplace:
- A. How would you describe the state of gender equality in your workplace during the COVID-19 response?
- B. Did you notice any differences in how male and female nurses were treated or expected to perform during the pandemic?

- 3. Awareness and Understanding of Gender Equality Policies:
- A. Were you aware of any gender equality policies implemented during the COVID-19 response?
  - B. How were these policies communicated to you and your colleagues?
  - C. In your view, were these policies effectively implemented?
  - 4. Impact of Policies on Work and Well-being:
- A. Did gender equality policies influence your work conditions (e.g., shift assignments, workload, decision-making involvement)?
  - B. How did these policies affect your physical and mental well-being, if at all?
- C. Can you describe any specific incidents where you felt supported or unsupported due to your gender during the pandemic?
  - 5. Challenges and Disparities:
- A. Were there any specific challenges you faced as a nurse during the pandemic that you believe were related to gender?
- B. Were caregiving responsibilities outside of work (e.g., family, children) considered or supported during the response phase?
  - 6. Institutional Support and Leadership
- A. How did leadership in your organization respond to gender-related concerns during COVID-19?
  - B. Do you feel your concerns or suggestions were heard and addressed?
  - 7. Suggestions for Improvement:
- A. What do you think could have been done differently to promote gender equality more effectively during the COVID-19 response?
- B. What recommendations would you give to health institutions for future emergency responses regarding gender equality?
  - 8. Closing Question:
- A. Is there anything else you would like to share about your experience related to gender equality during the COVID-19 response?

## Data Analysis:

At the end of the interviews, the participants were asked to speak freely in case of other points regarding the interview. All the interviews were recorded using a smartphone after taking the participants' permission and were transcribed into written text immediately. The data were analyzed using content analysis through the conventional method and Graneheim and

Lundman's five-step conventional content analysis approach. The methodology applied by Gernheim and Lundman for data analysis was comprised of a sequential series of five distinct steps. Firstly, following each interview, the transcriptions were meticulously scrutinized by reading them repeatedly, line by line and paragraph by paragraph, with the aim of obtaining a comprehensive comprehension. Secondly, the primary codes were extracted. Thirdly, based on similar codes, sub-categories, categories, and themes were formulated. Fourthly, the initial interview was categorized, and the subsequent interviews helped in finalizing the primary category. Finally, an endeavor was undertaken to ensure the utmost homogeneity within categories and the most remarkable heterogeneity between categories (13).

Graneheim (2017) has expressed ,"If the aim is to describe participants' experiences of ordinary phenomena and data is concrete and close to the participants' lived experience, it may be wise to limit the analysis to categories at a descriptive level. If the analytic process continues too far, the results can become so abstract and general that they could fit into any context, and thereby say nothing about the participants' unique experience in situation" (13).

The preliminary codes were categorized and labeled based on their conceptual similarity (subcategories). The sub-categories were compared and placed under main categories, which were more abstract (categories). The main categories were categorized under a more abstract concept (theme).

## Rigor and reliable data:

In order to ascertain the integrity of the data, four fundamental criteria—credibility, dependability, confirmability, and transferability—were employed, as delineated by Lincoln and Guba (14). The investigator engaged in an extended sojourn at the various research sites, which facilitated the establishment of rapport with the participants and a deeper comprehension of the contextual factors at play. The participants' feedback was additionally employed to authenticate the data and coding; subsequent to the coding process, the transcripts of the interviews were returned to the participants to confirm the fidelity of the codes and their corresponding interpretations. Moreover, the participants endeavored to represent a broad spectrum of diversity in terms of age, gender, experience, educational attainment, and professional settings. The supervisors meticulously examined the transcripts derived from the interviews. In conjunction with the supervisors, the derived codes and classifications were presented to several individuals well-versed in the nursing domain and qualitative research methodologies; these were subjected to thorough scrutiny. A high level of consensus was observed regarding the results extracted from the analysis. Ultimately, the categories and

excerpts from the interviews were translated into English by a qualified translator and subsequently refined by a professional editor.

### **Ethical considerations**

This study protocol was approved by the Ethics Committee of Semnan University of Medical Sciences (approval code: IR.SEMUMS.REC.1399.187). They were coordinated with Universities of Medical Sciences before sampling. The study objectives and procedure were also explained to the participants, and informed consent was obtained.


### **Results:**

The study was conducted with a cohort of fourteen nurses who were working specifically in COVID-19 units, six of whom were identified as male. The initial analysis of the interviews led to the acquisition of 175 codes. The participants' responses to the research question were organized into a main theme and its associated categories. Ultimately, the main theme was identified as "gender inequality in management policies and resource allocation," with four categories labeled as "inefficient management," "inequality in interactions," " Inequality in the management of available resources," and "inequality in consideration of staff needs" (see Table 2).

**Table 2:** Nurses' perspectives on gender equality policies during the pandemics in response phase

Main theme: Gender inequality in the management policies and resources allocation				
Categories	Sub-categories	Concepts		
Inefficient management	<ul> <li>Managerial absence</li> <li>Conflicts of rules and practices</li> </ul>	<ul> <li>Diminishing presence of field managers</li> <li>Reduced responsiveness to challenges facing wards</li> <li>Failure to comply with a statute for allocating working hours based on work experience, marital status, gender, and the number of children</li> <li>Absence of a nurse of the same gender to provide exceptional care to patients</li> <li>Imbalance in volume/type of work and nurses' gender and abilities</li> </ul>		
Inequality in Interactions	<ul> <li>Gender-based planning in professional interactions</li> <li>Proportional interaction requirements</li> </ul>	<ul> <li>Respect and the role of gender in administrative interactions</li> <li>Nurses' privacy</li> <li>Increasing caring relations and empathy in a crisis</li> <li>Interactions based on an understanding of needs</li> </ul>		
Inequality in the management of available resources	<ul><li>Organizing PPE by gender</li><li>Justice in resource allocation</li></ul>	<ul> <li>Medical and protective requirements based on gender</li> <li>Facilitated access to PPE</li> <li>Gender-appropriate protective cover</li> </ul>		
Inequality in consideration of staff needs	<ul> <li>Paying attention to staff demands</li> <li>Importance of paying attention to psychological needs</li> <li>Importance of paying attention to personal needs</li> <li>Importance of paying attention to gender needs</li> </ul>	<ul> <li>Paying attention to gender-appropriate emotional needs</li> <li>Paying attention to individual differences</li> <li>Paying attention to physiological differences</li> <li>Paying attention to gender-appropriate physiological needs</li> </ul>		

The conceptual model of the acquired data is illustrated in Figure 1.



Figur 1- Conceptual Model: Gender Inequality in Management Policies and Resource Allocation

This framework may serve for the subsequent purposes:

- Qualitative research on gender justice in healthcare management
- Policy analysis and development to promote equitable work environments
- Organizational audits focused on gender-sensitive planning and resource distribution
- Educational tools for nursing administration and gender studies

The main categories extracted in the present study, along with examples of the participants' quotations related to the concepts, were as follows:

## 1. Inefficient management:

- 1.1. Managerial absence
- 1.2. Conflicts of rules and practices

As subcategories, the first category identified was inefficient management, which was derived from managerial avoidance and conflicts arising from incongruent rules and practices. Furthermore, the related concepts encompassed the dwindling presence of field managers, diminished responsiveness to challenges encountered by wards, non-adherence to a statute

mandating the allocation of working hours based on work experience, marital status, and the number of children, the unavailability of a nurse of the same gender to provide exceptional care to patients, and a disparity in workload volume/type and nurses' gender and abilities.

In Iran, the provision of nursing care for female patients is predominantly carried out by female nurses, a practice shaped by prevailing cultural norms. Consequently, in cases where the female patient population surpasses the male patient population within the inpatient department and the female nurse's counterpart is a male nurse, it results in an escalation of workload for the female nurse.

When female nurses were engaged in extended and frequent shifts, particularly those who were married and had a spouse and children, their attention towards their patients was diminished as a result of the substantial workload and obligations. Numerous statements articulated by the study participants with regards to the aforementioned concepts were as follows:

- a. Concerning the concept of diminishing presence of field managers, one of the participants stated that: "I can almost say that the presence of some managers, in particular those who do most office work, not the ones with whom we have direct contacts, was deficient during this crisis, that is, practically neither senior supervisors nor the ones, whom I saw a lot before, like the infection control supervisors, who used to hang out and their main jobs should be now training other supervisors and generally the senior manager, was not seen in the COVID-19 unit during this pandemic at all." (27 years old, female)
- b. One of the participants reiterated the reduced responsiveness to challenges facing wards: "Due to the critical situation that occurred, we faced lots of problems in the wards. In these cases, we often used to call the supervisors, but unfortunately, the cooperation was not what it should have been." (27 years old, male)
  - Another nurse also stated, "Most nursing managers only insist that the work should be done anyway. Once the number of patients was very high and the workload was too high, they used to say that all work must be done anyway, and there was no way. They were adding fuel to the fire. However, at least it would have been better for them to give us a hand and get help from nurses from more private wards or be present in the ward themselves." (27 years old, female)
- c. In regards to failure to comply with a statute for allocating working hours based on work experience, marital status, and the number of children, one of the participants added that: "During this time, shifts were very long, but the important thing was that it

could become challenging for me if I was married, and I could not take care of the patients properly when I was on a shift." (30 years old, female)

One other Participant also said: "Female co-workers who had small children had similar shifts to ours, but they were constantly worried and even cried during shifts, and we had to do their jobs as well." (32 years old, male)

d. In regards to the absence of a nurse of the same gender to provide exceptional care to patients, one of the participants stated that: "In our ward, as a COVID-19 unit, the patients' families were not allowed to accompany them to help with some work. During most shifts, my co-workers and I were both men, and most of our patients were women, so we had hard times doing some procedures, such as bladder catheterization and intramuscular injection of drugs. The supervisors also did not help us." (28 years old, male)

Another nurse additionally maintained: "For example, one day, my co-worker was male. We divided the patients according to the care volume because we had to do so. My co-workers' patients were women, so I had to do some of their care, and my workload multiplied." (30 years old, female)

## 2. Inequality in Interactions:

- 2.1. Gender-based planning in professional interactions
- 2.2. Proportional interaction requirements

The second category of the study was centred on the topic of "Inequality in interactions". This was derived from the proportional interaction requirements and gender-based planning in professional interactions. The category encompassed various concepts such as respect and the role of gender in administrative interactions, privacy of nurses, enhancing caring relations and empathy during a crisis, and interactions based on an understanding of needs. Within the context of Iran, female nurses exhibit a tendency to engage in regulated interactions with male nurses and other individuals, influenced by religious considerations. Consequently, instances where their privacy is disregarded lead to feelings of insecurity. The study participants provided several statements based on the aforementioned concepts:

a. Regarding respect and the role of gender in administrative interactions, one of the participants noted that: "Our cooperation with different people was based on their moral characteristics in terms of gender. With a female co-worker, especially in terms of age, particularly when we were the same age, we could have a better and closer relationship." (27 years old, female)

- As well another participant said that: "We were usually more comfortable with coworkers, who had the same gender, but we also had no problem with male counterparts, and we used to try to maintain respect at the time of crisis." (37 years old, female)
- b. In regards to of nurses' privacy, one of the participants noted: "There should always be a framework for relationships, but in any case, my security should never be violated." (30 years old, female)
  - Moreover, another participant said that: "During the COVID-19 crisis, we were all under much stress, but we would not like anyone to invade our privacy, and that was annoying." (28 years old, male)
- c. One Participant added in regards to increasing caring relations and empathy in a crisis: "The difference between the COVID-19 crisis and normalcy was that our relationships became better and friendlier. The managers also became more intimate with us, and this helped us get through the difficult situation better." (28 years old, male)

  As well another participant said that: "It was true that the number of patients increased and we were terrified of getting sick, but some nursing managers and supervisors became more intimate with us, even by talking on the phone, and showed more compassion, which was good and inspiring." (37 years old, female)
- d. In regards to interactions based on an understanding of needs, a nurse stated that: "I have a small child, and my parents are also sick, so I am always worried about transmitting the disease to them in shifts, and I do not focus on the care I provide, but someday Ms. Akbari (one of the supervisors) came to our ward and found out that I was distraught, so she talked to me and reassured me, and that was very good." (34 years old, female)

## 3. Inequality in management of available resources:

- 3.1. Organizing PPE (personal protective equipment) by gender
- 3.2. Justice in resource allocation

The third category was related to the inequality in management of available resources, mined from organizing personal protective equipment by gender and justice in resource allocation, including concepts such as medical and protective requirements based on gender, facilitated access to personal protective equipment, and gender-appropriate protective covers. Some statements provided by the participants based on the given concepts were as follows:

a. Considering the concept of medical and protective requirements based on gender, most participants in this study stated that: "Men care less about personal protection coverage

than women, and they are more courageous in providing clinical care in COVID-19 wards." (37 years old, female)

Men's valiant actions in tending to individuals afflicted with COVID-19 may pose a risk to their own well-being; nevertheless, it enhances their proficiency in attending to covid-19 patients.

One of the nurses also said, "At the onset of this crisis, due to severe shortages of personal protective equipment, they were rationed in the wards ,and the head nurses were responsible for distributing the equipment between the nurses and other members of the treatment teams. Due to the cover we had on, in addition to the personal protective equipment, they got wet very quickly because of excessive sweating and had to be replaced sooner, but the number of our equipment was equal to that of men's." (30 years old, female)

- b. In regards to facilitated access to protective equipment, one Participant added that: "The head nurses, who were male, were much better at distributing protective equipment among the staff and better managing their shortages, but female head nurses and managers were more sympathetic to the staff and nurses. However, access to personal protective equipment was initially our biggest concern during the COVID-19 pandemic." (32 years old, male)
- c. Regarding the concept of gender-appropriate protective covers, one of the participants stated, "Our covers were the same during the COVID-19 crisis because personal protective clothing was very similar, and there was no difference between men and women. Of course, we hardly knew each other, and communication was difficult." (27 years old, male)

# 4. Inequality in consideration of staff needs:

- 4.1. Paying attention to staff demands
- 4.2. Importance of paying attention to psychological needs
- 4.3. Importance of paying attention to personal needs
- 4.4. Importance of paying attention to gender needs

The fourth category focused on the topic of disparity in fulfilling staff needs. It was drawn from considering the requirements of staff, their psychological needs, and their gender-specific necessities. The gender-related demands included a range of ideas like suitable emotional and physiological needs based on gender, and unique distinctions. The participants articulated several statements based on these aforementioned concepts, which included the following:

a. In regards to paying attention to gender-appropriate emotional needs, one Participant stated that: "I have often experienced that the authorities have been less strict with men and have had higher expectations of us as women, making us resentful with this gendered view." (30 years old, female)

The occupation of nursing aligns closely with the emotional and nurturing characteristics traditionally associated with women, leading nursing supervisors to potentially have higher expectations of female employees in the nursing profession. The establishment of such anticipations may appear justifiable as it pertains to the psychological attributes exhibited by female nurses. However, disparities arise in the workplace where both male and female nurses are deemed equivalent in their occupational duties and remuneration, thus necessitating a mutual enhancement of their favourable demeanour when tending to patients.

- b. In regards to paying attention to individual differences, one of the participants reiterated: "In my opinion, the main problem facing nursing managers was that they had the same expectations from everyone, and during this crisis, they only paid attention to the performance of nursing duties and care and did not consider nurses and their abilities at all." (27 years old, male)
- c. Regarding the concept of paying attention to physiological differences, a female nurse said, "Of course, I am not saying that women are less capable, but gender should be considered in the arrangement of shifts and wards. For example, I was on many shifts with a male co-worker; he used to help me a lot, did many activities that we could not do, and in general, it sometimes felt excellent to have a male co-worker in the ward." (30 years old, female)

### **Discussion:**

The COVID-19 pandemic has exposed several primary causes of inefficient management in nursing, significantly impacting healthcare delivery. Key factors include inadequate support for nursing staff, ethical dilemmas in resource allocation, and the psychological toll on healthcare professionals. These issues collectively hinder effective management and care provision during crises.

Inadequate Support for Nursing Staff

• Nurse managers often focused on logistical challenges, such as staffing and training, while neglecting the mental well-being of their teams(Botma & Roets, 2023).

• There was a critical shortage of personal protective equipment (PPE) and other resources, forcing nurses to make difficult ethical decisions regarding patient care(Gupta et al., 2024).

### **Ethical Dilemmas**

- The allocation of scarce resources, such as ventilators and ICU beds, raised ethical concerns about fairness and equity in patient care(Gupta et al., 2024).
- Nurses faced the challenge of balancing personal safety with professional obligations, leading to moral distress(Gupta et al., 2024).

## Psychological Impact

- The pandemic resulted in significant emotional strain on nurses, contributing to burnout and mental health issues(Aydogdu, 2022)(Hossny et al., 2022).
- Support systems, including mental health resources, were often insufficient, exacerbating the challenges faced by nursing staff(Dhaliwal et al., 2022).

Conversely, some argue that the pandemic has also fostered resilience and innovation in nursing management, prompting the development of new strategies and collaborative practices that may enhance future healthcare responses.

The present study was to explore gender equality policies from the perspectives of Iranian nurses working in pandemic situation. The study findings accordingly shed light on gender inequality in the management policies and resource allocation. Several groups were also developed, including inequality in management, inequality in interactions, inequality in management of available resources, and inequality in consideration of staff needs. In the meantime, some concepts such as conflicts of rules and practices, gender-based planning in professional interactions, organizing PPE by gender, and the importance of paying attention to psychological, personal, and gender needs, were extracted, which will be discussed below. Inefficient management was one of the issues affecting the nurses' views. Among these issues was the managerial absence; for example, reduced responsiveness to challenges facing the wards, as expressed by the interviewees. The managerial absence was also a new concept that had not been explored in nursing and other fields. Another issue was the diminishing presence of field managers. For example, the study participants mentioned the absence of the infection control supervisor in the wards for direct control and training. In their study, Hesselink et al. also reported that the presence and availability of nurse leaders during daily work were significant from the perspectives of nurses working in intensive care units (ICUs) (15). McSherry et al. also found that excellence in nursing care could occur if nursing managers and

leaders had responded to the complexities of reforms and changes (16), which was in line with the results of the present study.

The findings of the current study also highlighted a failure to adhere to a particular statute concerning the distribution of working hours, taking into account factors such as work experience, marital status, and the number of children. This serves as an illustration of conflicts arising from discrepancies between rules and actual practices. In this respect, Hoffman et al. had established that nurses working 12-hour shifts were younger, less experienced, and more stressed than their co-workers who were working 8-hour shifts. Once the differences in nursing experience had been controlled, similar stress levels in the nurses were observed (17).

Yu Liu et al., in their study in the nursing ward of a hospital, practically implemented sustainable support, dynamic human resource allocation, organization of pre-service training, monitoring of main stages of work, formulation of positive incentive methods, as well as the use of medical equipment, and had reported that such strategies had significantly boosted the capacity of the hospital nursing team to deal effectively with the COVID-19 pandemic As well, Wang et al. had concluded that the reasonable allocation of the nursing workforce along with other strategies could ensure safe, effective, timely, regular, and sustained centralized treatment of COVID-19 (18).

The imbalance in volume/type of work and nurses' gender and abilities also influenced their views towards policymaking in this study. Galbany-Estragués and Dolors had further shown that time pressure had separated clinical practices of nursing care from the context of nursing theory (19). Therefore, in case of a mismatch between workload and care with the nurses' gender and abilities, inefficient care and error would be observed. Randelle et al., in their study, had correspondingly revealed that a gender-balanced workforce, combined with unique care concepts, could strengthen the nursing profession and optimize abilities to provide customer services (20).

Interactions were also among the categories extracted in this study. In this line, Utoft highlighted three factors, viz. communication, leadership commitment, and holistic assessment, to improve gender equality in organizations (21), which was consistent with the present study.s findings' The concept of respect and the role of gender in administrative interactions were similarly clarified in the present study. In this regard, Zanielm et al. found that most nurses had mentioned gender as a significant factor in nurses' interactions, so women were more likely than men to experience interprofessional gender biases (22). However, Aguirre et al., examining gender patterns of interactions in an interprofessional health care

network, reported that network members were likelier to communicate with men and women without any preferences (23). This could be due to the cultural differences between the two study environments. Accordingly, in the study by Alghamdi et al., nurses, regardless of gender, showed job satisfaction and a greater understanding of transformational leadership style when their managers were male (24).

The study participants also reported increasing caring relations and empathy in crises, particularly in relationships between managers, staff, and co-workers. Unfortunately, no study was found assessing these two variables in relationships between nurses, so further research is suggested.

The management of available resources was another viewpoint among the nurses about policymaking during the COVID-19 pandemic. In this respect, Bambi et al. stated that long hours of using PPE were associated with adverse consequences such as discomfort, fatigue, and skin damage for most nurses working in COVID-19 ICUs. Nursing managers were thus required to adjust day and night shifts, taking into account the maximum duration of personal protective equipment use (25), which was in agreement with the findings of the present study. One of the concepts related to the management of available resources was the genderappropriate protective cover. The inequitable and disproportionate utilization of Personal Protective Equipment (PPE) emerged as a prominent indicator of gender disparity among nurses amidst the COVID-19 pandemic. Consequently, the presence of gender inequality persisted notwithstanding the equitable distribution of PPE, attributed to additional factors. No study related to this concept was found, but Ong et al. reported that using personal protective equipment could cause headaches or increase and exacerbate chronic ones in HCWs (26). Therefore, designing PPE based on the staff and gender helps improve their performance and maintain their physical health.

Although the consideration of staff needs in policymaking was acknowledged in this study, no case was found. Robert et al. had merely established that nurses were rarely involved in policymaking while implementing health policies (27), which could be why gender, as well as physical and psychological needs of nurses, had not been respected.

Although male nurses were less likely to directly report discrimination, 4 out of 6 male participants acknowledged an inequity in the distribution of high-risk shifts in their favor. This study pioneers in examining gendered dimensions of PPE distribution during COVID-19, revealing systemic inequity patterns in resource allocation despite quantitative parity - a critical gap unaddressed in prior literature."

## **Implications for International Contexts and Broader Health Systems**

The findings of this study, while rooted in a specific national context, have important implications for international health systems and global health policy. The gendered experiences of nurses during the COVID-19 response highlight the need for inclusive policy design that explicitly considers the diverse needs of healthcare workers based on gender, role, and social position. In broader health systems, this underscores the importance of embedding gender equity frameworks into emergency preparedness and workforce planning. Internationally, the study supports calls by organizations such as the WHO and UN Women to integrate gender-responsive approaches into health crisis management. The insights may also inform comparative policy development, guiding countries toward more equitable resource distribution, psychosocial support mechanisms, and staffing practices that reflect a commitment to gender justice in healthcare delivery.

# Strategies for Integrating Gender Equality into Pandemic Policy

## 1. Gender-Responsive Workforce Planning

- Ensure staffing policies account for gender-related roles, caregiving responsibilities, and occupational segregation (e.g., female-dominated nursing roles).
- Avoid disproportionate workloads for women through equitable shift scheduling and task distribution.

## 2. Inclusive Decision-Making

- Involve women—especially frontline health workers—in all levels of pandemic planning and policy-making committees.
- o Establish gender-balanced advisory boards and crisis management teams.

## 3. Sex-Disaggregated Data Collection & Analysis

- Collect and analyze data on infection rates, workload, PPE access, and psychological outcomes by gender to identify disparities in real-time.
- o Use these data to inform targeted interventions.

### 4. Equitable Resource Allocation

 Distribute personal protective equipment (PPE), rest spaces, and support services based on actual need, accounting for gender-specific requirements (e.g., fit and comfort of PPE).  Ensure pregnant and breastfeeding healthcare workers have appropriate accommodations and protections.

# 5. Supportive Work Policies

- Implement family-friendly policies (e.g., flexible hours, parental leave, childcare support) to reduce stress and burnout, especially among female health workers.
- Provide psychological support services tailored to gender-related stressors and experiences.

## 6. Anti-Discrimination and Harassment Protections

- Enforce zero-tolerance policies against workplace harassment, with particular attention to gender-based and sexual harassment during crisis conditions.
- 7. Create safe reporting channels and protections for whistleblowers.

# **Training and Awareness Programs**

- Integrate gender-sensitivity training into pandemic preparedness modules for managers and frontline staff.
- Build awareness of unconscious bias and its consequences in health crisis settings.

# 8. Accountability and Monitoring Mechanisms

- Establish systems to monitor the implementation of gender-sensitive measures throughout the pandemic lifecycle.
- Conduct regular audits and evaluations of gender equity outcomes in emergency response operations.

## **Study Limitations**

This study has several limitations that should be acknowledged. First, the transferability of the findings may be limited, as the experiences and perspectives of nurses are deeply embedded in specific organizational, cultural, and policy contexts. While efforts were made to include participants with diverse backgrounds, the findings may not be generalizable to nurses working in different healthcare systems or cultural settings. Second, the sample size was relatively small, involving 14 participants. Although this number is consistent with qualitative content analysis standards and saturation was achieved, a larger sample could have potentially uncovered additional nuances or divergent viewpoints, particularly across different roles or regions. Third, the study was conducted within a single-country context, which further limits

the applicability of the results to other national settings. Gender equality policies and their implementation during the COVID-19 pandemic likely varied across countries, influenced by broader sociopolitical and institutional structures. Therefore, the findings should be interpreted within the specific national and healthcare system framework in which the study was conducted. Future research should consider cross-country comparisons and include broader samples to enhance the understanding of how gender equality is addressed in healthcare crises globally.

### Conclusion

This study highlighted nurses' perceptions of gender inequality in management policies during the COVID-19 pandemic. Participants identified inefficient leadership, biased resource allocation, and unequal interactions as key barriers. These findings align with global evidence on crisis-driven gender disparities. To address this, we recommend:

Establishing gender-balanced decision-making teams, Regular audits of resource distribution, Training for managers on gender-sensitive leadership.

Future research should explore male nurses' perspectives and evaluate interventions to reduce inequities. Despite limitations (e.g., nurses' fatigue), this study underscores the urgent need for policy reforms before the next crisis."

The findings reveal that nurses' perception of managerial inadequacy and leadership gaps during crises significantly undermines workforce morale and erodes trust in institutional policies. Given the demonstrated impact of nursing staff's perspectives on policy formulation and crisis management efficacy, healthcare administrators must prioritize mechanisms for systematically integrating frontline nurses' insights into decision-making processes—particularly during emergency response planning.

To advance gender equity in nursing, we propose three critical action domains:

Root cause analysis: Rigorous mixed-methods investigations into structural and cultural determinants of occupational disparities between male and female nurses;

Intervention development: Design and implementation of evidence-based protocols to address identified inequities in resource allocation, leadership opportunities, and workplace interactions;

Impact evaluation: Longitudinal studies assessing the effectiveness of implemented solutions through metrics including retention rates, promotion equity, and staff satisfaction indices.

These recommendations carry particular urgency as healthcare systems worldwide prepare for future public health emergencies, where equitable workforce management will directly impact pandemic response effectiveness.

## **Limitations of the Study**

"This study has several limitations that should be acknowledged. First, the demanding workload and fatigue experienced by nurses during the pandemic crisis may have influenced their willingness to fully disclose sensitive experiences or critical perspectives regarding organizational policies. Some participants potentially self-censored their responses due to concerns about professional repercussions, despite researchers' efforts to create a safe and confidential interview environment. Second, the study's focus on a single healthcare setting during an extraordinary crisis period may limit the generalizability of findings to non-crisis situations or different healthcare systems. Finally, while data saturation was achieved with 14 participants, a larger sample size could potentially reveal additional dimensions of gender inequality experiences."

### **Conflicts of Interest**

"The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article."

# Acknowledgements

"We extend our deepest gratitude to the nurses who generously shared their time and experiences during this challenging pandemic period. Their invaluable contributions made this study possible. This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors."

### References

- 1. Sen NC, Chawla A. The Saga of Pandemics through the Literary Lens. Nusantara Journal of Behavioral and Social Science. 2023;2(4):105-10.
- 2. Mahalle PN, Ambritta PN, Sakhare SR, Kulkarni AP. Pandemic Problems. In: Mahalle PN, Ambritta PN, Sakhare SR, Kulkarni AP, editors. Foundations of Mathematical Modelling for Engineering Problem Solving. Singapore: Springer Nature Singapore; 2023. p. 143-55.
- 3. Rim D, Shin H, Jeon H, Lee Y-M, Shon S, Shim K, et al. Creating the 5A Nursing Response Model for the Emerging Pandemic: A Qualitative, Theory-Generating Study. Journal of Advanced Nursing.n/a(n/a).
- 4. Starr M, Webber-Ritchey KJ, Harris B, Simonovich SD. Exploring US Nursing Leadership During the Initial COVID-19 Pandemic Response: A Qualitative Descriptive Study to Guide Leadership Development for Future Emergent Situations. JONA: The Journal of Nursing Administration. 2024;54(2):118-25.
- 5. Mwenyango H. Impact of COVID-19 Pandemic on Women's Rights and Wellbeing: Analysis of the Ugandan Response to the Global Virus. Journal of Human Rights and Social Work. 2023;8(1):105-13.
- 6. Smith J, Davies SE, Feng H, Gan CCR, Grépin KA, Harman S, et al. More than a public health crisis: A feminist political economic analysis of COVID-19. Global Public Health. 2021;16(8-9):1364-80.
- 7. Ramakumar R, Eapen M. The legacy of public action and gender-sensitivity of the pandemic response in Kerala State, India. Economia Politica. 2022;39(1):271-301.
- 8. Smith JC. Representation in times of crisis: women's executive presence and gender-sensitive policy responses to crises. Journal of European Public Policy. 2023;30(10):1984-2009.
- 9. Essue BM, Kapiriri L, Mohamud H, Veléz MC, Kiwanuka S. Planning with a gender lens: A gender analysis of pandemic preparedness plans from eight countries in Africa. Health Policy OPEN. 2024;6:100113.
- 10. Ryan NE, El Ayadi AM. A call for a gender-responsive, intersectional approach to address COVID-19. Global Public Health. 2020;15(9):1404-12.
- 11. Callerstig A-C, Strid S. Gender mainstreaming in times of crisis: Missed opportunities in pandemic policymaking. Papers (Universitat Autònoma de Barcelona). 2023;108(3):0003-e3174.

- 12. Heidari S, Torreele E, Gülmezoglu AM, Sekalala S, Burke-Shyne N, Chappuis GL. A gender-responsive Pandemic Accord is needed for a healthier, equitable future. The Lancet. 2023;402(10418):2176-9.
- 13. Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. Nurse education today. 2004;24(2):105-12.
- 14. Polit DF, Beck CT. Nursing research: Generating and assessing evidence for nursing practice: Lippincott Williams & Wilkins; 2008.
- 15. Hesselink G, Branje F, Zegers M. What are the factors that influence job satisfaction of nurses working in the intensive care unit? A multicenter qualitative study. Journal of Nursing Management. 2023;2023(1):6674773.
- 16. McSherry R, Pearce P, Grimwood K, McSherry W. The pivotal role of nurse managers, leaders and educators in enabling excellence in nursing care. Journal of Nursing Management. 2012;20(1):7-19.
- 17. Hoffman AJ, Scott LD. Role stress and career satisfaction among registered nurses by work shift patterns. JONA: The Journal of Nursing Administration. 2003;33(6):337-42.
- 18. Wang H, Feng J, Shao L, Wei J, Wang X, Xu X, et al. Contingency management strategies of the Nursing Department in centralized rescue of patients with coronavirus disease 2019. International journal of nursing sciences. 2020;7(2):139-42.
- 19. Galbany-Estragués P, Comas-d'Argemir D. Care, autonomy, and gender in nursing practice: A historical study of nurses' experiences. Journal of Nursing Research. 2017;25(5):361-7.
- 20. Sasa RI, editor Male nurse: A concept analysis. Nursing forum; 2019: Wiley Online Library.
- 21. Utoft EH. Exploring linkages between organisational culture and gender equality work—an ethnography of a multinational engineering company. Evaluation and program planning. 2020;79:101791.
- 22. Zanielm MSc N, Manchanda M, Chary M, Nadeau R, Verstreken M, Shappell M, et al. The Role of Gender in Nurse-Resident Interactions: A mixed-methods study. 2020.
- 23. Aguirre-Duarte N, Carswell P, Kenealy T. Exploring gendered patterns of interaction within an inter-professional health care network. medRxiv. 2020.

- 24. Alghamdi MG, Topp R, AlYami MS. The effect of gender on transformational leadership and job satisfaction among Saudi nurses. Journal of advanced nursing. 2018;74(1):119-27.
- 25. Bambi S, Iozzo P, Lucchini A. New issues in nursing management during the COVID-19 pandemic in Italy. American Journal of Critical Care. 2020;29(4):e92-e3.
- 26. Ong JJ, Bharatendu C, Goh Y, Tang JZ, Sooi KW, Tan YL, et al. Headaches associated with personal protective equipment—A cross-sectional study among frontline healthcare workers during COVID-19. Headache: The Journal of Head and Face Pain. 2020;60(5):864-77.
- ra of C.

  Celline of the children of the child Anders RL, editor Engaging nurses in health policy in the era of COVID-19. Nursing 27.